

Edge Modes in the Hierarchical Fractional Quantum Hall Liquids with Coulomb Interaction

Takao Morinari and Naoto Nagaosa

Department of Applied Physics, University of Tokyo, Bunkyo-ku, Tokyo 113, Japan
(February 6, 2008)

Edge modes are studied for the hierarchical fractional quantum Hall liquids (FQHL) by treating the edge and bulk in a unified fashion. Within the RPA treatment of the composite boson effective theory with Coulomb interaction, one edge magneto-plasmon is shown to be the eigen-mode of the system. This mode decays algebraically perpendicular to the edge. All the other modes decay faster than power-law, and correspond to the neutral modes. However these neutral modes generally couple with the electromagnetic field within the magnetic length from the edge. The time-of-flight experiment in the $\nu = 2/3$ FQHL is discussed in terms of these results.

Keywords: D.electron-electron interactions, D. electronic transport, D. fractional quantum Hall effect

The fractional quantum Hall liquids(FQHL) are the incompressible quantum fluids, and the density fluctuation costs finite energy. Then the low lying excitations are possible only along the edge of the sample, which constitute the edge modes. These one-dimensional edge modes are described as the chiral-Tomonaga-Luttinger (CTL) liquids [1]. Recently, Milliken et al. observed at the Landau level filling factor $\nu = 1/3$, that the conductance G is proportional to $T^{2/\nu-2}$ which coincides with the prediction by Kane and Fisher about Tomonaga-Luttinger liquid [2]. Then it seems that the CTL theory works for the filling fraction $\nu = 1/(2k+1)$ (k : integer).

At the filling fraction $\nu = m/(mp+1)$ (m :integer, p :even), it is believed that FQHS has $|m|$ CTL modes. Especially in the case $m < 0$, it is predicted that there is one edge mode which has chirality opposite to the other modes.

However the time of flight experiment of the pulse injected into the $\nu = 2/3$ FQHL has detected only one chirality corresponding to the electron [3].

Kane et al. claimed that the backward scatterings caused by the impurities along the edge play a crucial role for the quantization of the Hall conductance [4]. These backward scatterings make the neutral mode massive, and only the charged mode carries the Hall current. However the fractional quantum Hall effect (FQHE) is a bulk phenomenon, and recently one of the present authors have shown that the Hall conductance is quantized *without* any randomness at the edge [5]. In this formalism, the edge and bulk are treated in a unified fashion and the edge modes are derived as the eigen-modes of the total system which are localized near the edge. With the short-range potential, the length scale perpendicular to the edge is of the order of the magnetic length $\ell_B = \sqrt{\hbar/eB}$. The edge currents, which are localized within this length, are driven by the electric field along the currents while the bulk currents are transverse.

On the other hand, a theory for the edge magneto-plasmon has been developed by Volkov and Mikhailov [6].

This theory takes into account the long-range Coulomb interaction but the relation between current density $\vec{J}(r)$ and the electric field $\vec{E}(r)$ is assumed to be that of the bulk up to the edge of the sample, i.e.,

$$J_\alpha(r) = \varepsilon_{\alpha\beta} \sigma_{xy} E_\beta(r). \quad (1)$$

where $\alpha, \beta = x, y$ and $\varepsilon_{\alpha\beta} = \pm 1$ is the antisymmetric tensor. Then this theory does not take into account the edge currents, and is considered to be relevant to the length scale much larger than ℓ_B , e.g., $\sim \mu\text{m}$.

Then the relationship between these two theories is not clear at present. Recently Orgad and Levit [7] have developed a theory for the edge mode for the FQHL with $\nu = 1/(2k+1)$. They take into account the long-range Coulomb interaction, which is similar in spirit to ref. [5] in the sense that the edge mode is derived from the bulk action. Orgad and Levit solved the RPA equations with the use of the ground state solution which is parametrized by the total charge of the system. They could treat also the case of the smooth change in the electron density near the edge. However their treatment is restricted to the case of $\nu = 1/(2k+1)$, i.e., the single mode case. On the other hand, Wen takes into account the long range nature of the Coulomb interaction in his one-dimensional action [8]. He concluded that only the magneto-plasmon is charged while all the others are neutral. However his treatment of the Coulomb interaction is rather phenomenological, and the spatial dependence perpendicular to the edge could not be studied.

In this paper we study the edge modes in the hierarchical FQHL in terms of the bulk action. This treatment naturally interpolates between the theory of magneto-plasmon by Volkov-Mikhailov [6] and that of the edge modes by Wen [1]. This work corresponds to the generalization of the method developed by Orgad and Levit [7] to the hierarchical case with the filling factor $\nu = m/(mp+1)$ (m :integer, p :even). It is found that only one mode is decaying algebraically and has the dispersion

like $\omega_k \sim -k \ln k$. This is the edge magneto-plasmon, and corresponds to the fluctuation of the total charge near the edge. All the others decay faster than power-law, probably exponential decaying within the magnetic length ℓ_B , and appear neutral in the longer length scale. The hydrodynamic argument reveals that the dispersion of these modes are $\omega_k \propto k$. Then our microscopic study confirms the conclusions by Wen clarifying its range of validity. From these results, we also discuss the time-of-flight experiment of the injected charge to the $\nu = 2/3$ FQHL.

In the case $\nu = m/(mp + 1)$, we begin with the bo-

son Chern-Simons Ginzburg-Landau theory. To begin with, we transform electrons to the composite fermions with p -flux of the Chern-Simons gauge field a_μ attached. Then the electron problem becomes that of the integer quantum Hall system with $|m|$ -filled Landau levels for composite fermions [9]. Each composite fermion is replaced by a boson with a flux $\phi_0 = ch/e$ attached to it. This flux corresponds to the other Chern-Simons gauge fields $a_{I\mu}$ with $I = 1, 2, \dots, |m|$. By integrating over a_μ after shifting $a_{I\mu}$ to $a_{I\mu} - a_\mu$, we obtain the Lagrangian density as

$$\mathcal{L} = \sum_{I=1}^{|m|} \left[\rho_I (-\hbar \partial_t \theta_I - U + e a_{I0}) - \frac{\rho_I}{2M_I} \left(\hbar \nabla \theta_I + \frac{e}{c} \mathbf{A} + \frac{e}{c} \mathbf{a}_I \right)^2 - \frac{\hbar^2}{2M_I} \left(\nabla \rho_I^{\frac{1}{2}} \right)^2 + \lambda_I \left(\frac{\rho_b}{|m|} \rho_I - \frac{1}{2} \rho_I^2 \right) \right] - \frac{1}{2} \int d^2 \mathbf{r}' V(\mathbf{r} - \mathbf{r}') [\rho(\mathbf{r}, t) - \rho_b(\mathbf{r})] [\rho(\mathbf{r}', t) - \rho_b(\mathbf{r}')] + \frac{e}{2\phi_0} \nu' \sum_{I,J} K_{IJ} \epsilon^{\mu\nu\lambda} a_{I\mu} \partial_\nu a_{J\lambda}. \quad (2)$$

Where U is an external potential, $K_{IJ} = p - (p|m| - 1) \delta_{IJ}$, $\nu' = 1/(|m|p - 1)$, $\rho = \sum_I \rho_I$, and ρ_b is the background positive charge density. The magnetic field B is in the $-z$ direction. Each boson field is characterized by the mass M_I and the short range hard core interaction λ_I . Now we generalize the treatment by Orgad and Levit [7] to derive the edge modes from eq.(2). Taking the variational derivative with respect to ρ , θ_I , and $a_{I\mu}$, the mean field equations are obtained. Putting $\mathbf{v}_I = (\hbar \nabla \theta_I + e/c \mathbf{A} + e/c \mathbf{a}_I)/M_I$, these mean field equations are

$$-\hbar \partial_t \theta_I - U + e a_{I0} - \frac{M_I}{2} \mathbf{v}_I^2 + \frac{\hbar^2}{2M_I} \frac{1}{\sqrt{\rho_I}} \nabla^2 \sqrt{\rho_I} + \lambda_I \left(\frac{\rho_b}{|m|} - \rho_I \right) - \int d^2 \mathbf{r}' V(\mathbf{r} - \mathbf{r}') [\rho(\mathbf{r}', t) - \rho_b(\mathbf{r}')] = 0, \quad (3)$$

$$\rho_I = \sum_J K_{IJ} \left(\frac{\nu'}{h} M_J \nabla \times \mathbf{v}_J + \nu' \frac{B}{\phi_0} \right), \quad (4)$$

$$\rho_I \mathbf{v}_I \times \hat{e}_z - \frac{c\nu'}{h} \sum_J K_{IJ} \left[\partial_t \left(M_J \mathbf{v}_J - \hbar \nabla \theta_J - \frac{e}{c} \mathbf{A} \right) + \frac{e}{c} \nabla a_{J0} \right] = 0. \quad (5)$$

The equation of continuity is derived from these equations. The ground state is obtained by solving these equations, but its explicit form is not needed in the following consideration.

We consider a sample which spreads over the semi-infinite plane ($x \geq 0$). In this geometry, we have the translational invariance along y -direction, then ρ_I and v_{Iy} are the functions of x only and $v_{Ix} = 0$. We choose the Landau gauge for the gauge fields $a_{I\mu}$, i.e. $\mathbf{a}_I = (0, a_{Iy}(x))$ and $\mathbf{A} = (0, -Bx)$. In this gauge choice, we see that $\partial_y \theta_I$ and $\partial_t \theta_I$ are functions of x only and $\partial_x \theta_I \propto v_{Ix}$ is equal to 0. Then we can write $\theta_I = q'_I y - (\mu_I/\hbar)t$ with constant q'_I and μ_I .

From (4), the excess charge $q_I (= \int_0^\infty dx (\rho_b/|m| - \rho_I))$ of I component boson field is related to q'_I by $q_I = (\nu'/2\pi) \sum_J K_{IJ} q'_J$. The boson fields equilibrate each other, and the chemical potential for I th component boson is equal to each other, i.e., $\mu_I = \mu$ (I -independent).

Linearizing the mean field equations around the ground state solutions, we get the RPA equations :

$$\delta \rho_I - \frac{\nu'}{h} \sum_J K_{IJ} M_J (\partial_x \delta v_{Jy} - \partial_y \delta v_{Jx}) = 0, \quad (6)$$

$$\rho_I \delta v_{Iy} + v_{Iy} \delta \rho_I - \frac{c\nu'}{h} \sum_J K_{IJ} \left[M_J \partial_t \delta v_{Jx} + \frac{e}{c} \partial_x \delta a_{J0} \right] = 0, \quad (7)$$

$$\rho_I \delta v_{Ix} + \frac{c\nu'}{h} \sum_J K_{IJ} \left[M_J \partial_t \delta v_{Jy} - \hbar \partial_t \partial_y \delta \theta_J + \frac{e}{c} \partial_y \delta a_{J0} \right] = 0, \quad (8)$$

$$\hbar \partial_t \delta \theta_I = -M_I v_{Iy} \delta v_{Iy} + f(\rho_I, \delta \rho_I) + e \delta a_{I0} - \lambda_I \delta \rho_I - \int d^2 \mathbf{r}' V(\mathbf{r} - \mathbf{r}') \delta \rho(\mathbf{r}', t). \quad (9)$$

where f is given by $f(\rho, \delta\rho) = -(\hbar^2/4M_I)\rho^{-3/2}(\nabla^2\rho^{1/2}) \times \delta\rho + (\hbar^2/4M_I)\rho^{-1/2}\nabla^2(\rho^{-1/2}\delta\rho)$. We will show below that one solution to these equations is obtained from the ground state configuration. For this purpose, we consider the ground-state solution with the fixed total charge, i.e., $q = \sum_I q_I$. We differentiate this ground state configuration ρ_I, v_{Iy}, a_{I0} and θ_I with respect to the total charge q , and the derivatives are denoted by $\delta\rho_I^{(s)}$ etc. When taking the derivative we choose $dq_I = dq/|m|$, which means that the charge of each component oscillates in phase. By using these, we try the solution to eqs.(6)-(9) which has the following form.

$$\begin{cases} \delta\rho_I = \delta\rho_I^{(s)}(x) \cos(ky - \omega t), \\ \delta a_{I0} = \delta a_{I0}^{(s)}(x) \cos(ky - \omega t), \\ \delta v_{Iy} = \delta v_{Iy}^{(s)}(x) \cos(ky - \omega t), \\ \delta\theta_I = \frac{2\pi}{\nu'} \frac{1}{k} \sin(ky - \omega t). \end{cases} \quad (10)$$

From eq. (8), it is seen that δv_{Ix} is of the order of k or ω and we put $\delta v_{Ix} = 0$. We put eq.(10) and $\delta v_{Ix} = 0$ into the lhs of eqs.(6) - (8), and keep only the lowest order terms in k and ω . After that we obtain the very equations for $\delta\rho^{(s)}$ etc that coincide with the equations obtained by differentiating the ground-state equations by the total-charge q . Then it has been shown that $\delta\rho_I$ etc. in eq.(10) satisfy the eqs.(6)-(8). The last equation (9) gives the dispersion ω_k of this mode. In the last term in the rhs of eq.(9), the Coulomb interaction leads to the modified Bessel function $K_0(|k|x)$. This function can be approximated as $K_0(|k|x) \cong \log(2e^{-\gamma}/|k|x)$ (γ is the Euler constant), because the ground state charge density is localized in the region of the order of magnetic length ℓ_B perpendicular to the edge. Then the dispersion is

$$\hbar\omega_k = \frac{\nu}{2\pi} \left(\frac{1}{|m|} \sum_{J=1}^{|m|} \frac{\partial\mu}{\partial q_J} \right) k - \nu \frac{e^2}{\pi} k \log \frac{2e^{-\gamma}}{|k|}. \quad (11)$$

From eqs.(3)-(5) the asymptotic behavior of the ground state solution for $x \gg \ell_B$ can be obtained as $v_{Iy} \sim (2\nu'/h)(|m|/\rho_b)(qe^2/x)$, $\rho_I - \rho_b/|m| \sim -(2\nu'^2/h^2)(|m|/\rho_b)(\sum_J K_{IJ}M_J)(qe^2/x^2)$, and $\partial_x a_{I0} \sim 2qe/x$. Then the mode obtained above decays algebraically perpendicular to the edge, i.e., $\delta v_{Iy}^{(s)} \sim (2\nu'/h)(|m|/\rho_b)(e^2/x)$, $\delta\rho_I^{(s)} \sim -(2\nu'^2/h^2)(|m|/\rho_b)(\sum_J K_{IJ}M_J)(e^2/x^2)$ and $\partial_x \delta a_{I0}^{(s)} \sim (2e/x)$. Then this mode can be identified as the edge magneto-plasmon because (i) it corresponds to the oscillation to the total charge q , i.e., the charge of each boson oscillates in phase. (ii) The spatial extent of this charged mode into the bulk is much larger than the magnetic length ℓ_B , and is insensitive to the details of the edge and the hierarchy structure of FQHL. This enables the classical treatment as given by Volkov-Mikhailov [6].

Thus we have found one eigenmode of the system. The next problem is if there are other charged modes which

have the similar properties as that above, i.e., $\omega_k \propto k \ln k$ and the power-law decay. To see this we analyze RPA equations (6)-(9) in the asymptotic region $x \gg \ell_B$. We are interested in the solutions with the power law decay, i.e., $\delta\rho_I \sim r_I/x^\alpha \cos(ky - \omega t)$, $\delta v_{Iy} \sim s_I/x^\beta \cos(ky - \omega t)$, $\partial_x \delta a_{I0} \sim t_I/x^\gamma \cos(ky - \omega t)$, where r_I, s_I, t_I and α, β, γ are some unknown constants. Putting these forms into eqs.(6)-(9), we neglect the terms which are the order of k or ω and keep only the dominant terms in the region $x \gg \ell_B$. Then we obtain the exponents as $\alpha = 2, \beta = \gamma = 1$ and

$$r_I = -\frac{\nu'}{h} \sum_J K_{IJ} M_J s_J, \quad (12)$$

$$s_I = \frac{|m|}{\rho_b} \frac{e\nu'}{h} t_I, \quad (13)$$

$$t_I = -2e \int dx' \delta\rho(x'). \quad (14)$$

It should be noted that if the total charge $\int dx' \delta\rho(x')$ is finite, it coincides with the asymptotic behavior of the mode obtained above. From the uniqueness of the solution to the differential equations, it is concluded that the edge magneto-plasmon is single. All the other modes are neutral in the sense that $\int dx' \delta\rho(x') = 0$, and hence decay faster than any power-law, probably exponentially, within the magnetic length ℓ_B . This is because ℓ_B is the only length scale when the Coulomb interaction is irrelevant, and the edge theory with short range interaction [1] [5] is applicable. Note that this condition mentions only about the integral of the charge, and generally the charge *density* of each neutral mode is not zero within the length scale of ℓ_B from the edge. This leaves the possibility to detect the neutral modes in terms of the electromagnetic measurements if the probe is localized within ℓ_B from the edge.

Now we study the other edge modes more explicitly. For this purpose we resort to the hydrodynamic treatment regarding the FQHL as a completely incompressible fluid up to the edge [10]. This method is rather phenomenological and less exact, and can not reproduce the correct behavior perpendicular to the edge. However all the edge modes can be derived in a simple way. We start from the following equation which can be obtained from eq.(3) by taking the gradient.

$$M_I [\partial_t \mathbf{v}_I + (\mathbf{v}_I \cdot \nabla) \mathbf{v}_I] = \frac{\hbar}{\nu'} \sum_J K_{IJ}^{-1} \rho_J (\mathbf{v}_J - \mathbf{v}_I) \times \hat{e}_z - e \mathbf{E} - \frac{e}{c} \mathbf{v}_I \times \mathbf{B} + \nabla P_I, \quad (15)$$

where

$$P_I = \frac{\hbar}{2M_I} \frac{1}{\sqrt{\rho_I}} \nabla^2 \sqrt{\rho_I} + \lambda_I \left(\frac{\rho_b}{|m|} - \rho_I \right) - \int d^2 \mathbf{r}' V(\mathbf{r} - \mathbf{r}') [\rho(\mathbf{r}', t) - \rho_b(x')] \quad (16)$$

and $\hat{e}_z$ is unit vector of z -axis. Now the approximation of the complete incompressibility is introduced. Because the density fluctuation is zero [10], $\nabla \times \mathbf{v}_I = (e/M_I c) \nabla \times (\mathbf{A} + \mathbf{a}_I) = 0$. From this equation the velocity $\mathbf{v}_I$ is expressed by the scalar potential ϕ_I as $\mathbf{v}_I = \nabla \phi_I$. In addition $\nabla \cdot \mathbf{v}_I = \nabla^2 \phi_I = 0$ from the equation of continuity. Then each $\mathbf{v}_I$ is expressed as $\mathbf{v}_I = \phi_I^{(0)} \nabla e^{iky - |k|x - i\omega t}$ with some constant $\phi_I^{(0)}$. For $V(\mathbf{r})$, we take the Coulomb interaction: $V(\mathbf{r}) = e^2/|\mathbf{r}|$. Now we introduce an approximation for the pressure P_I . We divide P_I into two contributions. One is from the long range Coulomb interaction, which corresponds to the third term in the rhs of eq. (16). The other is the surface tension due to the short-range interaction which roughly corresponds to the first and second terms. We write the surface tension energy as $E_{S.T.}^{(I)} = (\lambda'_I/2)\rho_b^2 \int dy \xi_I^2(y)$ with ξ_I being the displacement of ρ_I at y and the pressure from it is given by $(P_I)_{S.T.} = (|m|/\rho_b)(\delta E_{S.T.}^{(I)}/\delta \xi_I)$. The Coulomb interaction is taken into account by the 3-dimensional Coulomb integration [10]. The displacement ξ_I is related to ρ_I by the equation of continuity at the boundary, then we have from (15):

$$\omega \sum_J \frac{eB}{c} K_{IJ}^{-1} \phi_J^{(0)} + k \lambda'_I \rho_b \phi_I^{(0)} + 2\rho_b e^2 k \log \frac{k_0}{|k|} \cdot \sum_J \phi_J^{(0)} = 0, \quad (17)$$

where we ignored the ω^2 -order term and k_0^{-1} is some short range cut off. Note that $K_{IJ}^{-1} = \nu' p - \nu' \delta_{IJ}$, and define $a_I = -(eB/c)\nu'\omega + \lambda'_I \rho_b k$, $b = (eB/c)p\nu'\omega + 2e^2 \rho_b k \log(k_0/|k|)$ and $A_{IJ} = b + a_I \delta_{IJ}$, eq.(17) becomes

$$\sum_J A_{IJ} \phi_J^{(0)} = 0. \quad (18)$$

Then the eigenvalues are obtained by $\det A = 0$. This equation becomes

$$\left(\frac{eB}{c} p \nu' \omega + 2e^2 \rho_b k \log \frac{k_0}{k} \right) \sum_{J=1}^{|m|} \frac{1}{\rho_b \lambda'_I k - \frac{eB}{c} \nu' \omega} + 1 = 0. \quad (19)$$

If the solution ω to eq.(19) is order of $k \ln k$, we neglect the terms of order k and obtain

$$\hbar \omega_k \sim -\nu \frac{e^2}{\pi} k \log \frac{k_0}{|k|}. \quad (20)$$

For the other solutions, it can be seen that $\omega \sim k$. Then $A_{IJ} \sim b$ and we get from eq.(18), $\sum_{J=1}^{|m|} \phi_J^{(0)} = 0$. Then we find these mode are neutral modes. Especially in the case of $\nu = 2/3$, the solutions to eq.(19) are

$$\hbar \omega_k \sim -\frac{2}{3} \frac{e^2}{\pi} k \log \frac{k_0}{|k|} \quad (21)$$

and

$$\hbar \omega_k \sim \frac{3}{2} \rho_b \ell_B^2 (\lambda'_1 + \lambda'_2) k. \quad (22)$$

The former mode is the edge magnetoplasmon and the latter mode is the neutral mode. The propagating directions of these modes are opposite to each other.

Lastly we discuss the time-of-flight experiment in the $\nu = 2/3$ FQHL in terms of the results obtained above. As discussed below eq.(14), the neutral mode is not “neutral” locally within the length scale of ℓ_B . If one can detect such a local charge density, the neutral mode propagating in the opposite direction should show up. However in the experimental setup in ref. [3], the charge is introduced to a wide region near the edge whose length scale is typically $\sim 1\mu m (\gg \ell_B)$. Hence the total charge $\int dx \delta \rho(x)$ integrated perpendicular to the edge is relevant, and only the edge magneto-plasmon should be detected as observed experimentally.

The authors thank S. Levit, X. G. Wen, M. Kohmoto, A. M. Finkel'stein for useful discussions.

-
- [1] X. G. Wen, Phys. Rev. Lett. **64**, 2206 (1990).
 - [2] F. P. Milliken, C. P. Umbach, and R. A. Webb, Solid State Commun. **97**, 309 (1996).
 - [3] R. C. Ashoori et al. Phys. Rev. B **45**, 3894 (1992).
 - [4] C. L. Kane, M. P. A. Fisher, and J. Polchinski, Phys. Rev. Lett. **72**, 4129 (1994).
 - [5] N. Nagaosa and M. Kohmoto, Phys. Rev. Lett. **75**, 4294 (1995).
 - [6] V. A. Volkov, and S. A. Mikhailov, Sov. Phys. JETP **67**, 1639 (1988).
 - [7] D. Orgad and S. Levit, cond-mat/9507026.
 - [8] X. G. Wen, cond-mat/9506066
 - [9] B. Blok and X. G. Wen, Phys. Rev. B **42**, 8133 (1990).
 - [10] S. Giovanazzi, L. Pitaevskii and S. Stringari, Phys. Rev. Lett. **72**, 3230 (1994).

